

Extra-axial cerebello pontine angle medulloblastoma: A rare site of tumor

Sir,

Medulloblastoma is a common tumor of the posterior fossa, representing 20–25% of all pediatric neoplasms.^[1] The tumor often occurs in the cerebellar vermis and at the apex of the fourth ventricle.^[1,2] There are only a few reported cases of cerebellopontine (CP)-angle medulloblastoma in the literature, with most being intra-axial. The extra-axial site of this tumor remains a rarity.^[1,3]

This 4-year-old girl presented with left hemicranial headache followed by facial asymmetry with deviation of angle of mouth for 1 month. There was no other significant history.

On clinical examination, higher intellectual functions were normal, both pupils were equal and reacting to light, visual acuity/visual fields were normal, fundus—no papilloedema, left lower motor neuron facial paresis, and left-sided sensory neural hearing loss, other cranial nerves normal. No stigmata of neurofibromatosis was noted. A computerised tomography (CT) scan of the brain showed contrast enhancing extra-axial lesion in the left CP angle centered around internal acoustic meatus [Figure 1]. CT bone window did not show enlargement of the internal acoustic meatus or hyperostosis [Figure 1]. Magnetic resonance imaging (MRI) of the brain showed CP angle lesion which was hypointense on T1W and hyperintense on T2W image [Figure 2]. The lesion was brilliantly enhancing with contrast, and no dural tail or canalicular component noticed [Figure 3]. She underwent left retromastoid craniectomy and total excision of the lesion. It was grayish, moderately vascular, and soft. There was a clear plane between the tumor and cerebellum, whereas it was adherent to dura and tent laterally. The HPE was confirmed as desmoplastic medulloblastoma [Figure 4] with the high MIB-1 labeling index and S-100 negativity.

Medulloblastoma usually occurs in inferior medullary velum in the midline.^[2] However rarely it may occur laterally in the cerebellar hemisphere in the pediatric and adult age group^[1,3-5] with most being intra-axial. The extra-axial site of this tumor remains a rarity.^[1,3] Origin of medulloblastoma may be either from germinal cells or their remnants situated at the end of the posterior medullary velum or from remnants of the external granular layer.^[3,6] Their development in the CPA may be from the remnants of the external granular layer in the cerebellar hemisphere, including the flocculus which faces the CP




Figure 1: CT scan brain plain with contrast shows left extra-axial contrast enhancing lesion




Figure 2: MRI of the brain showed the CP angle lesion which was hypointense on T1W and hyperintense on T2W image




Figure 3: MRI of the brain contrast study axial, sagittal, and coronal section shows contrast enhancing extra-axial lesion




Figure 4: HPE suggestive of desmoplastic medulloblastoma

angle.^[3,6] In the CP angle, medulloblastomas though fifth, sixth, and eighth cranial nerves are frequently involved, these nerves were spared in this patient.^[5,7] CP angle medulloblastomas are very rare with nearly 36 cases published in the literature^[1,3,5] of which only 10 are in adults.^[1,3,5,7] The lack of association with any cerebellar tissue and the extra-axial location of the tumor made our patient's case quite rare. However, they are likely under-reported owing to publication bias and must be considered in the differential diagnosis of extra-axial CP angle lesions.

*Goutham Cugati, Manish Singh, Nigel Peter Symss,
Anil Pande, Vasudevan Madabushi Chakravarthy,
Ravi Ramamurthi*

*Department of Neurosurgery, Voluntary Health Services,
Adyar, Chennai, Tamil Nadu, India*

Address for correspondence:

Dr. Manish Singh,
Department of Neurosurgery
Voluntary Health Services, Adyar,
Chennai - 600113, Tamil Nadu, India
E-mail: manish007singh@yahoo.com

REFERENCES

1. Fallah A, Banglawala SM, Provias J, Jha NK. Extra-axial

medulloblastoma in the cerebellopontine angle. *Can J Surg* 2009;52:E101-2.

2. Mohan M, Pande A, Vasudevan MC, Ramamurthi R. Pediatric medulloblastoma: A review of 67 cases at a single institute. *Asian J Neurosurgery* 2008;2:63-69.

3. Singh M, Cugati G, Symss NP, Pande A, Vasudevan MC, Ramamurthi R. Extra axial adult cerebellopontine angle medulloblastoma: An extremely rare site of tumor with metastasis. *Surg Neurol Int* 2011;2:25.

4. Available from: <http://www.surgicalneurologyint.com/text.asp?2011/2/1/25/77178> [Last Cited on 2011 Feb 26].

5. Hubbard JL, Scheithauer BW, Kispert DB, Carpenter SM, Wick MR, Law ER Jr. Adult cerebellar medulloblastomas: The Pathological radiological and clinical disease spectrum. *J Neurosurg* 1989;70:536-44.

6. Kumar R, Bhowmick U, Kalra SK, Mahapatra AK. Pediatric cerebellopontine angle medulloblastomas. *J Pediatr Neurosci* 2008;3:127-30.

7. Stevenson L, Echlim F. Nature and origin of some tumors of cerebellum: Medulloblastoma. *Arch Neurol Psychiatry* 1934;31:93-109.

8. Jaiswal AK, Mahapatra AK, Sharma MC. Cerebellopontine angle medulloblastoma. *J Clin Neurosci* 2004;11:42-5.

Access this article online

Quick Response Code:



Website:
www.ijmpo.org

DOI:
10.4103/0971-5851.89801

Conference Announcement

Round Table Meeting on

“Targeted therapy in Malignancy: Where Future is!”

on 12th Nov 2011 at the AIIMS, New Delhi.

Contact Persons

Dr. Lalit Kumar, Organizing Secretary, Email: lalitaiims@yahoo.com,
Jose Pulickal, Email: Jose.pulickal@ranbaxy.com