

Primary granulocytic sarcoma of lip – A rare extramedullary presentation of myeloid leukemia

Tamojit Chaudhuri,
Sayan Paul¹, Kirti Srivastava¹

Department of Radiotherapy,
Sanjay Gandhi Post-graduate
Institute of Medical Sciences,
¹Chhatrapati Shahuji Maharaj
Medical University, Lucknow,
Uttar Pradesh, India

Address for correspondence:

Dr. Tamojit Chaudhuri,
Department of Radiotherapy,
Sanjay Gandhi Postgraduate
Institute of Medical Sciences,
Lucknow, Uttar Pradesh, India.
E-mail: tamojit.cnmc@gmail.com

ABSTRACT

A sixty year old male presented with a swelling in the upper lip. On cytopathology, the patient was diagnosed as a case of granulocytic sarcoma. His bone marrow examination was unremarkable. Patient was then treated by radiotherapy alone 30 Gray in 15 fractions and achieved complete response. Eighteen months after follow up the patient is absolutely normal. To our knowledge this is the first ever reported case of granulocytic sarcoma of lip.

Key words: *Granulocytic sarcoma, myeloid leukemia, radiotherapy*

A 60-year-old male presented with history of gradually progressive swelling in right upper lip for 1 month. On examination, a non-tender, firm, lumpy swelling was noted in the right upper lip [Figure 1]. Fine needle aspiration cytology (FNAC) of the lump was suggestive of granulocytic sarcoma featuring scattered single large cells with high N:C ratio, and oval to round, irregular,



Figure 1: Pretreatment lesion involving right upper lip




Figure 2: Microphotograph of the cytological smear of fine needle aspirate from the lip swelling showing immature myeloid cells including blasts (May Grunwald Giemsa, x400)



Figure 3: Three months post-treatment picture showing complete disappearance of the upper lip lesion

Access this article online

Quick Response Code:



Website:
www.ijmpo.org

DOI:
10.4103/0971-5851.116212

concentrically placed nuclei with granular single to multiple nucleoli. The cells had scanty to moderate amount of basophilic cytoplasm. The background consisted of RBCs, neutrophils, and lymphocytes [Figure 2]. Bone marrow aspiration and biopsy showed no marrow involvement by leukemic cells. In tumor board, the patient was planned for local radiotherapy and systemic chemotherapy. He received 30 Gy in 15 fractions local radiation with Co-60; but due to financial constraints, he did not receive chemotherapy.

Eighteen months following radiotherapy, the patient is completely asymptomatic [Figure 3]. His bone marrow and peripheral blood picture is still absolutely normal. To our knowledge, this is the first ever reported case of granulocytic sarcoma of lip.

How to cite this article: Chaudhuri T, Paul S, Srivastava K. Primary granulocytic sarcoma of lip - A rare extramedullary presentation of myeloid leukemia. *Indian J Med Paediatr Oncol* 2013;34:126-7.
Source of Support: Nil, **Conflict of Interest:** None declared.

New features on the journal's website

Optimized content for mobile and hand-held devices

HTML pages have been optimized of mobile and other hand-held devices (such as iPad, Kindle, iPod) for faster browsing speed.

Click on [**Mobile Full text**] from Table of Contents page.

This is simple HTML version for faster download on mobiles (if viewed on desktop, it will be automatically redirected to full HTML version)

E-Pub for hand-held devices

EPUB is an open e-book standard recommended by The International Digital Publishing Forum which is designed for reflowable content i.e. the text display can be optimized for a particular display device.

Click on [**EPub**] from Table of Contents page.

There are various e-Pub readers such as for Windows: Digital Editions, OS X: Calibre/Bookworm, iPhone/iPod Touch/iPad: Stanza, and Linux: Calibre/Bookworm.

E-Book for desktop

One can also see the entire issue as printed here in a 'flip book' version on desktops.

Links are available from Current Issue as well as Archives pages.

Click on  View as eBook