

Effect of areca nut chewing and maximal mouth opening in schoolgoing children in Ahmedabad

Azizfatema Munawer Khan,

Megha S. Sheth,

Romsha R. Purohit

Physiotherapy Department of SBB
College, VS General Hospital,
Ahmedabad, Gujarat, India

ABSTRACT

Context: Areca nut is chewed by itself and in various scented preparations. Areca nut chewing is widely practiced in many parts of Asia, including India. Users often consider it harmless and report a sense of well-being, but evidence has shown that it is far from harmless and can have multiple oral health implications such as oral submucosal fibrosis posing difficulty in opening mouth and carcinogenesis. Studies in India have reported increasing prevalence of this habit among schoolgoing children. The objectives of the study were to find the effect of areca nut chewing on mouth opening, compare it with the children not having this habit, and to find correlation between maximal mouth opening (MMO) and months of areca nut chewing. **Aims:** The aim of this study is to find the effect of areca nut chewing on MMO in schoolgoing children in Ahmedabad. **Settings and Design:** An observational analytical study was conducted across various schools of Ahmedabad. **Subjects and Methods:** A total of eighty male students of 12–14-year-old were included in the study. Group A included children having the habit of eating areca nut for 6 months or more, and Group B had children who did not have the habit of areca nut chewing. Children who had just started eating for <6 months were excluded from the study. MMO was calculated as distance from the edge of the upper incisor teeth to the edge of the lower incisor teeth using a calibrated fiber ruler. **Statistical Analysis Used:** Statistical analysis was performed by SPSS software version 20.0, with level of significance set at 5%. **Results:** Mean and standard deviation of MMO for Group A was 3.69 ± 0.5 cm and for Group B was 4.46 ± 0.4 cm. Statistically significant difference was found using Mann–Whitney U-test with $U = 239.500$ and $P = 0.0001$. Pearson's coefficient $r = -0.623$ and $P = 0.0001$ showed moderate correlation between months of chewing and MMO. The mean duration of chewing was found to be 1.5 years. **Conclusion:** There is difference in MMO between children chewing areca nut and not chewing it with a moderate correlation between months of eating areca nut and MMO.

Key words: *Areca nut chewing, children, maximal mouth opening*

Address for correspondence:

Dr. Azizfatema Munawer Khan,
38 Tube Welex Near
Sungan Cinema, Sarkhej,
Ahmedabad, Gujarat, India.
E-mail: azizfatmakhan@yahoo.com

INTRODUCTION

Areca nut (supari) is chewed by itself and in various scented preparations. Areca nut chewing is widely practiced in many parts of Asia, including India.^[1] Studies in India have reported increasing prevalence of this habit among schoolgoing children.^[1-3] Krishna D *et al.* in their study on the prevalence and characteristics of areca nut, gutka, and tobacco among schoolchildren of rural areas in and around Gandhinagar found 33.33% and 22.28% of the prevalence among male and female students, respectively.^[1] Users often consider it harmless and report a sense of well-being, but evidence has shown that it is far from harmless and can have multiple oral health implications such as oral submucosal fibrosis (OSMF) posing difficulty in opening mouth and carcinogenesis.^[4]

The objectives of the study were to find the effect of areca nut chewing on mouth opening, compare it with

the children not having this habit, and to find correlation between maximal mouth opening (MMO) and months of areca nut chewing.

SUBJECTS AND METHODS

An observational analytical study was conducted in two municipal schools of Ahmedabad, among 12–14-year-old male schoolgoing children. Group A included forty children having the habit of eating areca nut for 6 months or more, and Group B included forty children who did not have the habit of areca nut chewing. Children who had just started eating for <6 months were excluded from the study. A total of eighty children were included using a convenient sampling design.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Khan AM, Sheth MS, Purohit RR. Effect of areca nut chewing and maximal mouth opening in schoolgoing children in Ahmedabad. *Indian J Med Paediatr Oncol* 2016;37:239-41.

Access this article online

Quick Response Code:



Website:
www.ijmpo.org

DOI:
10.4103/0971-5851.195734

Permission for data collection was sought from the principals of the schools. Oral informed consent of the children included in the study was taken. Demographic data were collected using a self-made questionnaire. MMO was measured. For measuring MMO, children were asked to open their mouth fully till no further opening was possible. Average of three readings was taken. Distance from the edge of the upper incisor teeth to the edge of the lower incisor teeth was measured. A calibrated ruler was used for measurement.^[5] Level of significance was kept at 5%.

Between-groups analysis was done using Mann–Whitney test. Correlation between months of eating and MMO was found using Spearman’s correlation analysis.

RESULTS

Mean MMO was compared for Group A and Group B; the difference in mean MMO for Group A and Group B was found to be statistically significant. Comparison of mean MMO and SD of Group A and Group B is shown in Table 1. Correlation between months of chewing and MMO was done using Spearman’s correlation as shown in Graph 1. A moderate negative correlation ($r = -0.623$) was found between months of areca nut chewing and MMO. The mean duration of areca nut was found to be 1.5 years.

DISCUSSION


The mean MMO was found to be significantly reduced in children who had the habit of chewing areca nut (3.69 cm) than those who did not have the habit (4.46 cm).

MMO for Indian boys is 4.6 cm, 4.8 cm, and 5.2 cm in age groups of 6–8, 8–10, and 10–12 years, respectively.^[5] It is

Table 1: Comparison of mean maximal mouth opening and standard deviation of Group A and Group B

Groups	Mean±SD	U	P
A	3.69±0.5	239.500	0.0001
B	4.46±0.4		

SD – Standard deviation



Graph 1: Correlation of months of areca nut chewing and maximal mouth opening

seen that as age increase mouth opening increases. The MMO in the present study was less than above findings even though age group was slightly more.

Furthermore, a moderate correlation was found between months of chewing areca nut and MMO. Reddy *et al.* in their study on correlation of the clinical grading of OSMF to various chewing habits found that the relative risk of OSMF increased with duration, frequency, and style of chewing habits for longer duration, which is similar to the present findings.^[6] Nigam *et al.* in their study on 1000 individuals having habit of chewing nut or gutka for 1 year or more found the prevalence of OSMF to be 6.3%. Out of which 58.7% individuals had reduced mouth opening with MMO of <3 cm and 38.09% had altered taste sensations.^[7]

Dere *et al.* in their study on the prevalence and characteristics of areca nut, gutka, and tobacco among schoolchildren of rural areas in and around Gandhinagar found 10% boys and 2% females to be suffering with OSMF with reduction in MMO.^[11] Gupta *et al.* in their study on series of two cases, one girl aged 11 years and eating areca nut for 7 years and other boy aged 10 years and eating areca nut since 6 years, showed that both had complaints of reduced mouth opening, discomfort, and burning sensation particularly when eating spicy food found MMO to be 1.4 cm and 1.3 cm, respectively.^[8] Reduced mouth opening was more than seen in the present study probably because duration of eating was maximum 3 years.

The major chemical constituents of areca nut are carbohydrates, fats, and alkaloids.^[9] Arecoline, one of the alkaloids, is the main agent for carcinogenic effect.^[10] Application of these substances to human fibroblasts induces fibroblastic proliferation and collagen production and thus progressive difficulty in mouth opening leading to OSMF.^[11] OSMF is a premalignant condition of oral cavity strongly associated with nut and gutka chewing. It is characterized by burning sensation of mouth especially while eating spicy food, fibrosis of oral soft tissue resulting in decreased mouth opening, and pallor of oral mucosa.^[12] The present study, however, did not evaluate the other above characteristics.

The malignant transformation rate of OSMF has been found to be 13%–40% worldwide, whereas it is 7.6% in the Indian population.^[13] Goel *et al.* studied OSMF by dividing 100 cases into three stages; Stage 1 - MMO >4.1 cm, Stage 2 - restricted MMO 2–4.4 cm, and Stage 3 MMO <2 cm. They found that 60% of individuals had restricted MMO falling into Stage 2.^[14]

It is likely that children who are regular chewers at early age will be dependent on the nut and continue the chewing habit in their adult life, which will make them more prone to OSMF, unless appropriate intervention is made.^[8]

Only boys were included in the study which is one of the limitations of the study.

The collected data can be used for the purpose of planning areca nut chewing control interventions and evaluation. Studies to see effect of exercises on MMO in OSMF need to be given proper emphasis.

CONCLUSION

There is difference in MMO between the children chewing areca nut and not chewing it. There is a correlation between months of eating areca nut and MMO.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Dere K, Choudhry P, Bhaskar V, Ganesh M, Venkataraghvan K, Shah S. Prevalence and characteristics of Areca nut, Gutka and Tobacco among school children of rural areas in and around Gandhinagar. *J Adv Oral Res* 2014;5:20-5.
2. Khandelwal A, Khandelwal V, Saha MK, Khandelwal S, Prasad S, Saha SG. Prevalence of areca nut chewing in the middle school-going children of Indore, India. *Contemp Clin Dent* 2012;3:155-7.
3. Sarfraz M, Ali A, Mirza T. Prevalence and characteristics of areca nut chewing habit among school going children in Karachi. *J Dow Univ Health Sci Karachi* 2014;8:111-16.
4. Karemore TV, Karemore VA. Etiopathogenesis and treatment of oral sub mucosal fibrosis. *J Indian Acad Oral Med Radiol* 2011;23:598-602.
5. Kumar A, Mehta F, Goel M, Dutta S, Hooda A. Maximal mouth opening in Indian children using a new method. *J Craniomaxillary Dis* 2012;1:79-85.
6. Reddy V, Anjari PV, Banda NR, Reddy P. Oral Sub mucous fibrosis: Correlation of clinical grading to various habit factors. *Int J Dent Clin* 2011;3:21-4.
7. Nigam NK, Aravinda K, Dhillon M, Gupta S, Reddy S, Srinivas Raju M. Prevalence of oral submucous fibrosis among habitual gutkha and areca nut chewers in Moradabad district. *J Oral Biol Craniofac Res* 2014;4:8-13.
8. Gupta VK, Malhotra S, Patil R, Tripathi A. Oral submucous fibrosis at pediatric age, now time to think: Series of two cases. *Indian J Med Paediatr Oncol* 2013;34:107-10.
9. Jayalakshmi A, Mathew G. The areca nut palm. Kerala: Central Plantation Crops Research Institute; 1982. p. 225-44.
10. Hoffman D, Hecht SS. Advances in tobacco carcinogenesis, smokeless tobacco and betel quid: Control of tobacco related cancers and other diseases. Bombay: Oxford University Press; 1992. p. 193-204.
11. Harvey W, Scutt A, Meghji S, Canniff JP. Stimulation of human buccal mucosa fibroblasts *in vitro* by betel-nut alkaloids. *Arch Oral Biol* 1986;31:45-9.
12. Murti PR, Bhonsle RB, Gupta PC, Daftary DK, Pindborg JJ, Mehta FS. Etiology of oral submucous fibrosis with special reference to the role of areca nut chewing. *J Oral Pathol Med* 1995;24:145-52.
13. Rajalalitha P, Vali S. Molecular pathogenesis of oral submucous fibrosis – A collagen metabolic disorder. *J Oral Pathol Med* 2005;34:321-8.
14. Goel S, Ahmed J, Singh MP, Nahar P. Oral sub mucous fibrosis, a clinico- histopathological comparative study in population of Southern Rajasthan. *J Carcinog Mutagen* 2010;1:1-4.