Cancer risk of general people due to using joss stick for religious worshiping

Sir,

Worshipping is the basic religious practice seen anywhere around the world. In Asian, the use of joss stick for worshipping is common, and burning of joss stick can cause smoke that contains several hazardous chemicals.^[1,2] The exposure is a great concern for the workers in worship places, shrine, and temples.^[3,4] It is noted that the long-term exposure might result in carcinogenesis. In a recent publication from Thailand, a country where joss stick is widely used for religious worshipping, it is estimated that the lifetime cancer rates of the worker at worship place due to exposure to benzene and ethylbenzene in joss stick smoke are equal to 1–345 and 0.3–19 cases per million.^[5] Nevertheless, the risk for general people has not been assessed. Basically, in Thailand, general people will visit worship place on a specific period, which is called "temple day," about 4 times per month, and the people will stay and worship at the temple about half an hour per time. Hence, the period of exposure for the general people is significantly less than that of temple workers who have to work in temple every day for about 12 h per day. Here, the authors try to estimate the cancer risk due to exposure to joss stick smoke in religious worshipping. Based on the previous data from Thailand, the estimated cancer risk can be calculated. The period of exposure for general people compared to the temple worker is estimated to be about 0.0056 ([0.5 h/12 h] × 4 day/30 day). Hence, the estimated calculated lifetime cancer rates of general people due to exposure to benzene and ethylbenzene in joss stick smoke are equal to 0.0056-1.9167 and 0.0017-0.1056 cases per million.^[5] Based on the observed rate, the risk is very low, and it might imply that the general worshiping practice of the local people should not be considered as a cancer risk behavior. In fact, this result can also support the previous report that addresses no respiratory problem due to joss stick use on.^[6]

Financial support and sponsorship Nil.

Conflicts of interest

There are no conflicts of interest.

Beuy Joob, Viroj Wiwanitkit¹

Sanitation 1 Medical Academic Center, Bangkok, Thailand, ¹Tropical Medicine, Hainan Medical University, Haikou, China. E-mail: beuyjoob@hotmail.com

REFERENCES

- 1. Lin TS, Shen FM. Trace metals in Chinese joss stick smoke. Bull Environ Contam Toxicol 2003;71:135-41.
- Chuang HC, BéruBé K, Lung SC, Bai KJ, Jones T. Investigation into the oxidative potential generated by the formation of particulate matter from incense combustion. J Hazard Mater 2013;244-245:142-50.
- Lung SC, Kao MC. Worshippers' exposure to particulate matter in two temples in Taiwan. J Air Waste Manag Assoc 2003;53:130-5.
- Kuo CY, Yang YH, Chao MR, Hu CW. The exposure of temple workers to polycyclic aromatic hydrocarbons. Sci Total Environ 2008;401:44-50.
- Maspat V. Health risk assessment of workers exposure to BTEX from Chinese smoke at worship places in Bangkok. Bangkok: Chulalongkorn University; 2013.
- Liou SH, Yang JL, Cheng SY, Lai FM. Respiratory symptoms and pulmonary function among wood dust-exposed joss stick workers. Int Arch Occup Environ Health 1996;68:154-60.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.



How to cite this article: Joob B, Wiwanitkit V. Cancer risk of general people due to using joss stick for religious worshiping. Indian J Med Paediatr Oncol 2016;37:307.