

Classical or pylorus-preserving pancreatoduodenectomy in pancreatic and periampullary cancer: "The jury is still out!"

Hüttner *et al.*^[1] present an update of their previous Cochrane review^[2] comparing pylorus-preserving pancreaticoduodenectomy (pylorus-preserving Whipple [PPW]) versus classical pancreaticoduodenectomy (classic Whipple [CW]) for surgical treatment of periampullary and pancreatic carcinoma. The updated review included two new trials^[3,4] published since the last version taking the tally of included trials to eight overall.^[3-11] The present review,^[1] similar to the previous one,^[2] looked at the primary outcomes including pancreatic fistula, delayed gastric emptying (DGE), and biliary leak rates, secondary outcomes such as survival and postoperative mortality, and perioperative parameters such as intraoperative blood loss, red blood cell transfusion, operating time, postoperative bleeding, wound infection, pulmonary complications, necessity for re-operation, duration of hospital stay, quality of life, and status of resection margin.

Table 1 compares the various outcomes analyzed in the two manuscripts. The findings of the updated review largely echo the previous results, namely, there exists no difference between the CW and PPW in terms of overall morbidity, mortality, and survival. However, a novel finding in the updated version is the significantly increased risk of DGE in patients undergoing a PPW. On the flipside, PPW

has been consistently shown to be associated with shorter operating times, lower intraoperative blood loss, and hence, a reduced need for blood transfusion.

As was lucidly stated by the authors in the discussion, the analysis does reveal problems related to heterogeneity between the studies, variation in definitions used within the trials as well as the process of randomization. Even the intention-to-treat analysis was specified in only one trial.^[10] Also, not considered in the analysis are the data on whether the patients received neoadjuvant or adjuvant therapy – a significant determinant or morbidity, mortality, and survival.

Despite its inherent shortcomings, this manuscript represents the most updated review published on the topic and provides a "real world" overview of the differences (or lack of them) between the procedures for the management of pancreatic and periampullary cancers considering that the authors in the Collaboration with the Cochrane system have conducted an analysis reducing the influence of bias and heterogeneity as best they could.

However, this review certainly provides sufficient justification for a large randomized controlled trial investigating CW and PPW in pancreatic and periampullary cancers to more conclusively answer this question.

Table 1: Comparison of the outcomes analyzed between the two studies

Outcome	Diener <i>et al.</i> , 2014 ^[2]			Hüttner <i>et al.</i> , 2016 ^[1]		
	Number of studies	Effect size	Inference	Number of studies	Effect size	Inference
Primary outcome						
Pancreatic fistula	5	OR 0.86 (0.41, 1.81)	No difference	7	OR 0.95 (0.49, 1.84)	No difference
Delayed gastric emptying	5	OR 2.35 (0.72, 7.61)	No difference owing to heterogeneity	7	OR 3.03 (1.05, 8.70)	Favored CW
Biliary leak	3	OR 1.35 (0.10, 18.55)	No difference	5	OR 0.96 (0.15, 6.17)	No difference
Secondary outcome						
Survival	3	HR 0.84 (0.61, 1.16)	No difference	3	HR 0.84 (0.61, 1.16)	No difference
Postoperative mortality	5	OR 0.49 (0.17, 1.40)	No difference	7	OR 0.64 (0.26, 1.54)	No difference
Perioperative parameters						
Intraoperative blood loss	1	MD -0.76 (-0.96, -0.56)	Favors PPW	5	MD -0.32 (-0.62, -0.03)	Favors PPW
Red blood cell transfusion	2	MD -0.65 (-1.92, 0.61)	No difference	5	MD -0.47 (-0.86, -0.07)	Favors PPW
Operating time	3	MD -68.26 (-105.70, -30.83)	Favors PPW	7	MD -45.22 (-74.67, -15.78)	Favors PPW
Postoperative bleeding	3	OR 0.74 (0.29, 1.88)	No difference	5	OR 0.74 (0.32, 1.74)	No difference
Wound infection	4	OR 0.85 (0.35, 2.05)	No difference	4	OR 0.85 (0.35, 2.05)	No difference
Pulmonary complications	3	OR 0.67 (0.29, 1.58)	No difference	3	OR 0.67 (0.29, 1.58)	No difference
Necessity for re-operation	2	OR 0.82 (0.38, 1.75)	No difference	3	OR 0.80 (0.38, 1.68)	No difference
Duration of hospital stay	1	MD -1.80 (-8.94, 5.34)	No difference	5	MD 0.26 (-2.04, 2.56)	No difference
Status of Ro resection margin	-	-	No difference	4	OR 0.92 (0.39, 2.15)	No difference

OR – Odds ratio; HR – Hazard ratio; MD – Mean difference; CW – Classical Whipple; PPW – Pylorus-preserving Whipple

Savio George Barreto

*Department of Gastrointestinal Surgery, Gastrointestinal Oncology, and Bariatric Surgery, Medanta Institute of Digestive and Hepatobiliary Sciences, Medanta - The Medicity, Sector 38, Gurgaon, Haryana, India
E-mail: georgebarreto@yahoo.com*

REFERENCES

- Hüttner FJ, Fitzmaurice C, Schwarzer G, Seiler CM, Antes G, Büchler MW, *et al.* Pylorus-preserving pancreaticoduodenectomy (pp Whipple) versus pancreaticoduodenectomy (classic Whipple) for surgical treatment of periampullary and pancreatic carcinoma. *Cochrane Database Syst Rev* 2016;2:CD006053.
- Diener MK, Fitzmaurice C, Schwarzer G, Seiler CM, Hüttner FJ, Antes G, *et al.* Pylorus-preserving pancreaticoduodenectomy (pp Whipple) versus pancreaticoduodenectomy (classic Whipple) for surgical treatment of periampullary and pancreatic carcinoma. *Cochrane Database Syst Rev* 2014;11:CD006053.
- Srinarmwong C, Luechakietisak P, Prasitvilai W. Standard Whipple's operation versus pylorus preserving pancreaticoduodenectomy: A randomized controlled trial study. *J Med Assoc Thai* 2008;91:693-8.
- Taher MA, Khan ZR, Chowdhury MM, Nur-E-Elahi M, Chowdhury AK, Faruque MS, *et al.* Pylorus preserving pancreaticoduodenectomy vs. standard Whipple's procedure in case of carcinoma head of the pancreas and periampullary carcinoma. *Mymensingh Med J* 2015;24:319-25.
- Bloechle C, Broering D, Latuske C. Prospective randomized study to evaluate quality of life after partial pancreatoduodenectomy according to Whipple versus pylorus preserving pancreatoduodenectomy according to Longmire-Traverso for periampullary carcinoma. *Dtsch Gesellschaft Chir* 1999;Suppl 1:661-4.
- Lin PW, Lin YJ. Prospective randomized comparison between pylorus-preserving and standard pancreaticoduodenectomy. *Br J Surg* 1999;86:603-7.
- Lin PW, Shan YS, Lin YJ, Hung CJ. Pancreaticoduodenectomy for pancreatic head cancer: PPPD versus Whipple procedure. *Hepatogastroenterology* 2005;52:1601-4.
- Seiler CA, Wagner M, Bachmann T, Redaelli CA, Schmied B, Uhl W, *et al.* Randomized clinical trial of pylorus-preserving duodenopancreatectomy versus classical Whipple resection-long term results. *Br J Surg* 2005;92:547-56.
- Paquet KJ. Comparison of Whipple's pancreaticoduodenectomy with the pylorus-preserving pancreaticoduodenectomy – A prospectively controlled, randomized long-term trial [Vergleich der partiellen Duodenopancreatektomie (Whipple-Operation) mit der pylorus-erhaltenden Zephaloduodenopancreatektomie – Eine prospektive kontrollierte, randomisierte Langzeitstudie]. *Chir Gastroenterol* 1998;14:54-8.
- Tran KT, Smeenk HG, van Eijck CH, Kazemier G, Hop WC, Greve JW, *et al.* Pylorus preserving pancreaticoduodenectomy versus standard Whipple procedure: A prospective, randomized, multicenter analysis of 170 patients with pancreatic and periampullary tumors. *Ann Surg* 2004;240:738-45.
- Wenger FA, Jacobi CA, Haubold K, Zieren HU, Müller JM. Gastrointestinal quality of life after duodenopancreatectomy in pancreatic carcinoma. Preliminary results of a prospective randomized study: Pancreatoduodenectomy or pylorus-preserving pancreatoduodenectomy. *Chirurg* 1999;70:1454-9.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Access this article online

Quick Response Code:



Website:
www.ijmpo.org

DOI:
10.4103/0971-5851.195747

How to cite this article: Barreto SG. Classical or pylorus-preserving pancreatoduodenectomy in pancreatic and periampullary cancer: "The jury is still out!". *Indian J Med Paediatr Oncol* 2016;37:209-10.