

## Malignancy-Associated Microangiopathic Hemolytic Anemia and Thrombocytopenia

Sir,

Abdulla *et al.* nicely described two Indian patients who presented with microangiopathic hemolytic anemia (MAHA) and thrombocytopenia due to disseminated malignancy (DM).<sup>[1]</sup> I presume that underlying human immunodeficiency virus (HIV) infection might contribute to the revelation of DM in the studied two patients. That contribution could be addressed in dual aspects. On the one hand, it is obvious that patients with HIV infection have increased tendency to have various neoplastic lesions compared to healthy individuals. The increased propensity of neoplasms among HIV-positive patients has been thought to be related to different factors, including coinfection with oncogenic viruses, immunosuppression, and life prolongation secondary to the use of antiretroviral therapy.<sup>[2]</sup> To the best of my knowledge, HIV infection is a significant health hazard in India. The available data pointed out 0.26% HIV seroprevalence compared with a global average of 0.2%.<sup>[3]</sup> On the other hand, thrombotic microangiopathy is associated with HIV infection<sup>[4]</sup> and it could be the first clinical manifestation of HIV infection.<sup>[5]</sup> I presume that implementing the diagnostic workup of viral overload and CD4 count estimations was solicited in the studied two patients. If that diagnostic workup was accomplished and it revealed underlying HIV infection, the two cases in question could be truly regarded novel case reports. This is because concurrent HIV infection and DM uncovered by MAHA and thrombocytopenia have never been reported in the literature so far.

### Financial support and sponsorship

Nil.

### Conflicts of interest

There are no conflicts of interest.

**Mahmood Dhahir Al-Mendalawi**

*Department of Paediatrics, Al-Kindy College of Medicine, University of Baghdad, Baghdad, Iraq*

*Address for correspondence:*

*Prof. Mahmood Dhahir Al-Mendalawi,  
P. O. Box: 55302, Baghdad Post Office, Baghdad, Iraq.  
E-mail: mdalmendalawi@yahoo.com*

**Submitted:** 11-May-2018

**Accepted in Revised Form:** 21-Jun-2018

**Published:** 17-Feb-2020

### References

1. Abdulla MC, Alungal J, Nair LK, Musambil M. Malignancy associated microangiopathic hemolytic anemia and thrombocytopenia. *Indian J Med Paediatr Oncol* 2018;39:254-6.
2. Valencia Ortega ME. Malignancies and infection due to the human immunodeficiency virus. Are these emerging diseases? *Rev Clin Esp* 2018;218:149-55.
3. Paranjape RS, Challacombe SJ. HIV/AIDS in India: An overview of the Indian epidemic. *Oral Dis* 2016;22 Suppl 1:10-4.
4. Gilardin L, Malak S, Schoindre Y, Galicier L, Veyradier A, Coppo P. Human immunodeficiency virus-associated thrombotic microangiopathies. *Rev Med Interne* 2012;33:259-64.
5. Sarmiento M, Balcells ME, Ramirez P. Thrombotic microangiopathy as first manifestation of acute human immunodeficiency virus infection: A case report and review of the literature. *J Med Case Rep* 2016;10:152.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

### Access this article online

Quick Response Code:



Website:

www.ijmpo.org

DOI:

10.4103/ijmpo.ijmpo\_111\_18

**How to cite this article:** Al-Mendalawi MD. Malignancy-associated microangiopathic hemolytic anemia and thrombocytopenia. *Indian J Med Paediatr Oncol* 2019;40:594.

© 2020 Indian Journal of Medical and Paediatric Oncology | Published by Wolters Kluwer - Medknow