Original Article

Clinicopathological Profile of Head and Neck Squamous Cell Carcinoma

Abstract

Background: Head and neck squamous cell carcinoma (HNSCC) constitutes one of the most common malignancies in the world. The geographic location influences the etiologic factors and site of tumor. Aims and Objectives: The present study was carried out to illustrate the clinicopathologic profiles of HNSCC patients since data regarding these tumors from eastern region of India are scarce. Materials and Methods: A prospective study was undertaken for 2 years in which patients with histologically proven HNSCC were included. The clinicopathologic features of each case were analyzed. Results: A total of 108 cases were included in the study, among which 79 (73.15%) were male and 29 (26.85%) were female. Mean age of the patients was $53.21 (\pm 12.17)$ years. The most common risk factor was smoking (63 cases, 58.33%) followed by tobacco or betel nut chewing (41 cases, 37.96%). The common patterns of presentation included ulcerated lesion (51 cases, 47.22%), whitish lesion (28 cases, 25.93%), and hoarseness of voice (11 cases, 10.19%). The most common sites involved were buccal mucosa (36 cases, 33.33%) and dorsal surface of the tongue (26 cases, 24.07%). The most common site for exophytic tumors was buccal mucosa (9 out of 23, 39.13%) and that for ulceroproliferative lesions was tongue (9 out of 17, 52.04%). Microscopically, well-differentiated (Grade I) tumors were most common (67 cases, 62.04%) followed by moderately differentiated (Grade II) tumors (38 cases, 35.19%). A statistically significant correlation was obtained between anatomic site and grade of the tumor. Conclusion: Patients of HNSCC from the eastern region of India have distinctive features with regard to macroscopic appearance and microscopic grade of their tumors.

Keywords: Anatomical distribution, demographic pattern, head and neck squamous cell carcinoma, histopathological grade, risk factors

Introduction

Head and neck cancers (HNCs) are the sixth most common malignancy in the world. Majority of HNCs are squamous cell carcinoma (HNSCC), and among these, oral cancers account for up to 40% of all malignancies.^[1] France, India, and Brazil have the highest age-adjusted rates of HNC and Indian females account for the highest age-adjusted rates of HNC in the world.^[2,3] Anatomical sites affected by HNC varies worldwide. In India, the most common site is oral cavity, whereas pharynx accounts for majority of cases in France.^[4] reflecting the influence of exposure to different risk factors in the population.^[2,5] Because of indigenous practice of chewing pan and betel leaf with tobacco, cancers of tongue as well as buccal mucosa are frequent.^[6] Prognosis of oral squamous cell carcinoma (OSCC) is influenced by tumor primary site, nodal involvement,

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms. tumor thickness, and status of the surgical margins and is greatly influenced by the stage of the disease, especially pathological TNM (pTNM).^[7]

A literature search revealed that data regarding distribution in different age groups and sex, anatomic distribution, and clinicopathologic profile of HNSCC are scarce in the eastern region of India. The present study aims to report these features from a series of HNSCC from a referral center in Kolkata, West Bengal.

Materials and Methods

The duration of the present study was 2 years. All patients presented with HNC were screened for histologically proven SCC. These cases were evaluated for inclusion in the present study. Patients having other variants of malignancy, recurrence, metastatic carcinoma from unknown primary site, and lesions diagnosed only by cytological methods were excluded from the study group.

How to cite this article: Dasgupta S, Chakrabarti S, Deb AR. Clinicopathological profile of head and neck squamous cell carcinoma. Indian J Med Paediatr Oncol 2019;40:369-73.

Senjuti Dasgupta, Sudipta Chakrabarti¹, Asit Ranjan Deb²

Departments of Pathology and ²Radiotherapy, Medical College, ¹Department of Pathology, ESI PGIMSR, Manicktala, Kolkata, West Bengal, India

Submitted: 14-Sep-2017 Revised: 18-Apr-2018 Accepted: 21-Jun-2018 Published: 04-Dec-2019

Address for correspondence: Prof. Sudipta Chakrabarti, Department of Pathology, ESI PGIMSR, Manicktala, Kolkata, West Bengal, India. E-mail: sudiptach@gmail.com



For reprints contact: reprints@medknow.com

Data regarding sex, age, duration of illness, risk factors, clinical presentation of the lesion, location, and size were obtained. Clinical presentation of the lesions was classified in three groups: ulcers (including plain ulcers, and exophytic ulcerated masses), leukoerythroplakias, and tumors presenting both ulcers and leukoerythroplakic areas.^[8] Anatomic location of tumor in the oral cavity was subclassified based on the following sites: buccal mucosa (including buccal sulcus/mucobuccal fold), alveolar mucosa and gingiva (including retromolar area), border of tongue, floor of mouth (with extension to ventral tongue), upper and lower lips, soft palate, and tonsil areas. Histological slides were reviewed for the classification of tumors according to the following grade: well-differentiated (WD), moderately differentiated (MD), and poorly differentiated (PD) tumors according to the WHO criteria.^[9] In cases where the tumor was excised by radical surgery, pTNM was performed. Standard statistical methods were applied to analyze these data.

Results

A total of 123 cases were initially evaluated for inclusion in the study. After applying the exclusion criteria, 108 cases were selected as the study population. The anatomic distribution of cases is shown in Table 1. Buccal mucosa is the most common region (36 cases, 33.33%) followed by dorsal surface of tongue (26 cases, 24.07%) [Table 1].

79 Male accounted for cases (73.15%) while 29 cases (26.85%) occurred in female (M:F - 2.7:1) [Table 2]. Majority of anatomic locations had a sex ratio in favor of males, the highest being buccal mucosa (M:F - 8:1). Exceptionally, cases of neoplasms arising from alveolar mucosa/gingiva/retromolar area had a sex ratio in favor of females (M:F - 0.45:1). The age range of cases in the study group noted was between 31 and 81 years. Most cases occurred in the age group of 40-59 years (58 cases, 53.70%) [Chart 1]. Mean age of all cases was 53.21 (±12.17) years [Table 2]. However, cases of SCC arising from pharynx had a mean age of 63.22 (±7.40) years.



Chart 1: Distribution of cases according to age range

Consumption of alcohol was noted in 45 (41.67%) cases. No addition of these substances was noted in 21 cases (19.44%). Clinically, the most common presentation was ulcerated lesion (51 cases, 47.22%) followed by whitish lesion or mass (28 cases, 25.93%) and hoarseness of voice (11 cases, 10.19%). On examination, anemia was noted in 29 (26.85%) cases and palpable regional lymph node was detected in 18 (16.67%) cases. Clinically, duration of symptoms in most cases (60.19%)

was <6 months and only 10.19% cases had duration of >1 year [Table 3]. Most common clinical appearance

History of smoking was obtained in 63 (58.33%) cases

and tobacco/beetle nut chewing in 41 (37.96%) cases.

Table 1: Distribution of cases according	rding to anatomic			
location				
Location	Number of cases (%)			
Buccal mucosa	36 (33.33)			
Dorsal surface of tongue	26 (24.07)			
Alveolar mucosa/gingiva/retromolar area	16 (14.81)			
Larynx	13 (12.04)			
Pharynx	9 (8.33)			
Floor of mouth	3 (2.78)			
Ventral surface of tongue	1 (0.93)			
Palate	1 (0.93)			
Nasal cavity	2 (1.85)			
Lower lip	1 (0.93)			
Total	108 (100.00)			

Table 2: Demographic data of the study group according to anatomical site

to unatomical site				
Site	Average age	Male:female		
Buccal mucosa	50.08±11.39	8:1		
Dorsal surface of tongue	50.62±13.20	1.6:1		
Alveolar mucosa/	54.31±11.97	0.45:1		
gingiva/retromolar area				
Larynx	53.69±12.76	All male		
Pharynx	63.22±7.40	3.5:1		
Others	61.75±7.40	2:1		
Total	53.23±12.17	2.7:1		

Table 3: Clinical characteristics of cases in the study
group (<i>n</i> =108)

Parameters	Number of cases (%)
Duration of symptoms	
<6 months	65 (60.19)
6-12 months	32 (29.63)
>12 months	11 (10.19)
Predominant clinical appearance of tumors	
Ulcer	30 (27.78)
Ulcer with leukoerythroplakia	43 (39.81)
Exophytic mass	28 (25.93)
Leukoerythroplakia	7 (6.51)

of tumors was ulcer with leukoerythroplakia (39.81%) followed by ulcerated lesions (27.78%).

Specimen of radical surgery was obtained in 45 cases (41.67%) and biopsy of the tumor was available in 63 cases (58.33%) for pathological study [Table 4]. Microscopically, Grade I (WD) were the most common (67 cases, 62.04%) followed by Grade II (MD) tumors (38 cases, 35.19%), while Grade III (PD) tumors were the least common.

Additional pathological observation was documented in specimens of radical surgeries (n = 45). Most tumors had the greatest dimension between 2 and 4 cm (18 cases, 40.00%). Mean tumor thickness was 0.98 (±69) cm. Tumor depth was <1 cm in 17 cases (37.78%) and 1–2 cm in

Table 4: Histological characteristics of tumors		
	Number of cases (%)	
Tumor differentiation (<i>n</i> =108)		
Grade I well differentiated	67 (62.04)	
Grade II moderately differentiated	38 (35.19)	
Grade III poorly differentiated	3 (2.78)	
Specimens of radical surgery (<i>n</i> =45)		
Greatest dimension of tumor		
2 cm or less	12 (26.67)	
2-4 cm	18 (40.00)	
>2 cm	15 (33.33)	
Tumor depth		
<1 cm	17 (37.78)	
1-2 cm	16 (35.56)	
>2 cm	12 (26.67)	
Status of surgical resection margins		
Involved	16 (35.56)	
Uninvolved	29 (64.44)	
Lymph node metastasis		
Involved	10 (22.22)	
Not involved	35 (77.78)	
Lymphovascular invasion	7 (15.56)	
Perineural invasion	5 (11.11)	
Bone/cartilage invasion	2 (4.44)	
Dysplasia at margin	12 (26.67)	

16 cases (35.56%). Surgical resection margin was involved by the tumor in 16 (35.56%) cases, while metastatic deposit in the lymph nodes was seen in 10 (22.22%) patients. Other notable findings were lymphovascular invasion (7 cases, 15.56%), perineurial invasion (5 cases, 11.11%), and bone or cartilage invasion (2 cases, 4.44%).

In 62 cases (57.41%) documented, macroscopic appearance of tumors was available (as noted either in resected specimens or by imaging) [Table 5]. While exophytic lesions were most common in the buccal mucosa (9 out of 23, 39.13%), ulceroproliferative lesions were most common in the tongue (9 out of 17, 52.04%). However, microscopic examination revealed that Grade I SCC is the most common histological grade in both locations (72.22% and 65.38%).

Chi-square test was used to find a correlation between sets of two variables, i.e., microscopic grade of tumor to age of patient, site of tumor, duration in months, and macroscopic appearance of lesion. The correlation between microscopic grade and anatomic site of tumor was found to be statistically significant [Table 6].

Discussion

OSCC is the major bulk of tumor constituting HNSCC as well as oral cancers. There is a wide variation in the incidence of OSCC in different regions of the world. However, high proportion of occurrence of OSCC among these groups represents the exposure to different carcinogenic substances and their genetic predisposition.

Among the known risk factors, smoking, chewing tobacco, betel quid, areca nuts, and drinking alcohols are significant. In a case–control study conducted in the US involving 1114 patients of oral and pharyngeal cancer, it was observed that risks of cancer increased in a multiplicative manner in persons who are both smokers and drinkers. There were similar relative risk patterns among different races and sexes.^[10] Similarly, in a large study involving 5458 cases of HNC in Andhra Pradesh, India, only 21.22% cases were free from such risk factors.^[11] In the present study, only 19.44% cases had no addiction of these

Table 5: Macroscopic appearance and microscopic grade of tumors according to site of the lesion							
Tumor features	Buccal	Dorsal surface	Alveolar mucosa/gingiva/	Larynx	Pharynx	Others	Total
	mucosa	of Tongue	retromolar area/lower lip				
Macroscopic appearance (<i>n</i> =62)	23	17	10	7	3	2	62
Ulcer	6	3	5	1	1	2	18
Ulceroproliferative	7	9	0	6	2	0	24
Exophytic	9	3	4	0	0	0	16
Infiltrative	1	2	1	0	0	0	4
Total							
Microscopic grade (<i>n</i> =108)	36	26	16	13	9	8	108
Grade I	26	17	11	3	5	5	67
Grade II	10	9	5	9	3	2	38
Grade III	0	0	1	1	1	0	3

correlation of grade and anatomic site (P<0.05)				
Parameter		Microscopic grade		P
	Grade I (<i>n</i> =67)	Grade II (n=38)	Grade III (n=3)	
Age (years)				
<50	27	13	1	0.814778
50 or more	40	25	2	
Anatomic site				
Oral cavity	56	25	1	0.026487
Other sites	11	13	2	
Duration (months)				
<6	41	23	1	0.627198.
>6	26	15	2	
Macroscopic appearance				
Ulcer/ulcer with leukoerythroplakia	46	25	2	0.954941
Other lesions	21	13	1	

Table 6: Correlation between histopathological grade with clinical features. Chi-square test is significant in the
correlation of grade and anatomic site ($P < 0.05$)

causative substances. Other risk factors include ultraviolet radiation and poor oral hygiene.^[12] Human papillomavirus is now emerging as a significant risk factor in nonsmokers and in cancers involving tonsil, oropharyngeal region, and tongue.^[13]

In worldwide, most studies had documented that HNSCC mostly occurred in males.^[8,10,11] The male predominance is also apparent in all age groups and in all anatomic sites.^[8,11] A similar trend has been observed in the present study where M:F ratio was 2.7:1 with the highest ratio in buccal mucosa (M:F - 8:1). These patterns of sex ratio may be explained by the fact that males are more commonly addicted to risk factors such as alcohol, smoking, and chewing tobacco which are consumed concurrently.^[8,14]

Age groups that have been affected most commonly in the present study are 40-59 years (58 cases, 53.70%) followed by 60-79 years (33 cases, 30.56%). No case has been detected in persons below 30 years in the present study. The preponderance of such age distribution may be explained by the fact that these persons are usually exposed to risk factors for a sufficient duration, resulting in neoplastic changes. The occurrence of OSCC below 40 years of age is uncommon and ranges from 0.4% to 6%.^[8,15]

In the present study, the most common site affected was buccal mucosa (33.33%) followed by dorsal surface of tongue (24.07%). Addala et al. reported similar sites of affection among patients of HNC in Andhra Pradesh, India.^[11] The predilection for buccal mucosa in patients from Southeast Asia has been attributed to areca nut- and tobacco-chewing habits of people in this region.^[16] In fact, the variations in the commonly affected sites in different geographic locations are guided by the etiologic factors which are prominent in that particular zone. In American and European countries, the most common site affected is the border of the tongue, while the lower and upper gingiva have been reported to be commonly affected in Nigeria.^[8]

It was found in the current study that duration of symptoms in most cases (60.19%) was <6 months. In only 10.19% of cases, the duration exceeded 1 year. A comparable time span has been reported by Shenoi et al. in their study. They attributed the delay in diagnosis to low literacy rate, poor financial status, and tendency of Indian patients to resort to home remedies before seeking medical help.^[17] Interestingly, Pires et al. observed that the mean time of complaint before diagnosis was longer in females than in males^[8]

Many studies report that the site of predilection in HNC is apparently influenced by gender though exact reasons justifying such distribution remain obscure. The palate, alveolar mucosa, buccal mucosa, and gingiva are more commonly affected in females, whereas in males, border of tongue, ventral surface of tongue, and floor of mouth are the more common sites.^[8,14] In the present study, alveolar mucosa, gingiva, and retromolar area were found to be the common sites of affection in females as compared to buccal mucosa and dorsal surface of tongue, which showed a predilection for males.

Varied clinical appearances of tumors were appreciated in the present study, the most common being ulcer with leukoerythroplakia (39.81%), followed by ulcerated lesions (27.78%). Pires et al. however noted in their study that ulcers were the most common presentation (62%). They reported ulcers with leukoerythroplakia in 21% cases only.^[8]

In the present study, microscopically, Grade I (WD) tumors were the most common (67 cases, 62.04%) followed by Grade II (MD) tumors (38 cases, 35.19%). Most studies report HNC to be histologically WD or MD tumors.^[8] However, Effiom et al. in their study reported PD tumors to be the most common.^[18] Pires et al. found a relationship between the grade of tumor and gender of the patients. Males were mostly diagnosed with MD and PD tumors while females with WD and MD tumors. They

also suggested an association of grade of tumor with site. According to their observations, WD tumors are more common in buccal mucosa, buccal sulcus, and lower lip, whereas borders of tongue, ventral tongue, floor of mouth, and gingiva are the common sites for MD tumors.^[8] In the present study, Grade I SCC was most common in both buccal mucosa and tongue (72.22% and 65.38%, respectively).

The correlation between microscopic grade and anatomic site of the tumor was found to be statistically significant in the present study. An attempt was made by Shenoi *et al.* to correlate sets of two variables, i.e., site and habits, staging and anatomic site, stage and duration of illness, stage and habits, and stage and age of patient. However, none of these correlations was reported to be statistically significant by them.^[17]

Conclusion

The present study has revealed that males having an age range of 40–59 years were the most common population affected by HNSCC. Such tumors are strongly associated with addiction and only 19.44% cases were not addicted to any substances. Buccal mucosa is the most common anatomic location of HNSCC in the present study followed by dorsal surface of tongue. Ulceroproliferative and exophytic lesions were the most common macroscopic appearances while Grade I (WD) SCC was the predominant microscopic pattern in most sites. The histologic grading was statistically significant when correlated with anatomic site. Although these features have similarities with other studies conducted in India and abroad, some distinctive observations noted in this study were the macroscopic appearance and microscopic grade.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

- 1. Vokes EE, Weichselbaum RR, Lippman SM, Hong WK. Head and neck cancer. N Engl J Med 1993;328:184-94.
- Sankaranarayanan R, Masuyer E, Swaminathan R, Ferlay J, Whelan S. Head and neck cancer: A global perspective on epidemiology and prognosis. Anticancer Res 1998;18:4779-86.
- Parkin DM, Whelan SL, Ferlay J, Teppo L, Thomas DB, editors. Cancer Incidence in Five Continents. IARC Scientific Publisheds No. 155. Vol. 8. Lyon: IARC; 2002.

- Coleman M, Estève J, Damiecki P, Arslan A, Renard H. Trends in Cancer Incidence and Mortality. IARC Scientific Publisheds 121. Lyon: International Agency for Research on Cancer; 1993. p. 27-38.
- Menvielle G, Luce D, Goldberg P, Bugel I, Leclerc A. Smoking, alcohol drinking and cancer risk for various sites of the larynx and hypopharynx. A case-control study in France. Eur J Cancer Prev 2004;13:165-72.
- Thomas G, Hashibe M, Jacob BJ, Ramadas K, Mathew B, Sankaranarayanan R, *et al.* Risk factors for multiple oral premalignant lesions. Int J Cancer 2003;107:285-91.
- Noguchi M, Kinjyo H, Kohama GI, Nakamori K. Invasive front in oral squamous cell carcinoma: Image and flow cytometric analysis with clinicopathologic correlation. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2002;93:682-7.
- Pires FR, Ramos AB, Oliveira JB, Tavares AS, Luz PS, Santos TC, *et al.* Oral squamous cell carcinoma: Clinicopathological features from 346 cases from a single oral pathology service during an 8-year period. J Appl Oral Sci 2013;21:460-7.
- Barnes L, Eveson JW, Reichart P, Sidransky D. World Health Organization Classification of Tumors – Pathology & Genetics – Head and Neck Tumors. Lyon: IARC Press; 2005.
- Blot WJ, McLaughlin JK, Winn DM, Austin DF, Greenberg RS, Preston-Martin S, *et al.* Smoking and drinking in relation to oral and pharyngeal cancer. Cancer Res 1988;48:3282-7.
- Addala L, Pentapati CK, Reddy Thavanati PK, Anjaneyulu V, Sadhnani MD. Risk factor profiles of head and neck cancer patients of Andhra Pradesh, India. Indian J Cancer 2012;49:215-9.
- Guha N, Boffetta P, Wünsch Filho V, Eluf Neto J, Shangina O, Zaridze D, *et al.* Oral health and risk of squamous cell carcinoma of the head and neck and esophagus: Results of two multicentric case-control studies. Am J Epidemiol 2007;166:1159-73.
- Kreimer AR, Clifford GM, Boyle P, Franceschi S. Human papillomavirus types in head and neck squamous cell carcinomas worldwide: A systematic review. Cancer Epidemiol Biomarkers Prev 2005;14:467-75.
- 14. Kruse AL, Bredell M, Grätz KW. Oral cancer in men and women: Are there differences? Oral Maxillofac Surg 2011;15:51-5.
- Llewellyn CD, Johnson NW, Warnakulasuriya KA. Risk factors for squamous cell carcinoma of the oral cavity in young people – A comprehensive literature review. Oral Oncol 2001;37:401-18.
- Johnson NW, Jayasekara P, Amarasinghe AA. Squamous cell carcinoma and precursor lesions of the oral cavity: Epidemiology and aetiology. Periodontol 2000 2011;57:19-37.
- Shenoi R, Devrukhkar V, Chaudhuri, Sharma BK, Sapre SB, Chikhale A, *et al.* Demographic and clinical profile of oral squamous cell carcinoma patients: A retrospective study. Indian J Cancer 2012;49:21-6.
- Effiom OA, Adeyemo WL, Omitola OG, Ajayi OF, Emmanuel MM, Gbotolorun OM, *et al.* Oral squamous cell carcinoma: A clinicopathologic review of 233 cases in Lagos, Nigeria. J Oral Maxillofac Surg 2008;66:1595-9.